

Author Index

Aoyama, K. 87
Áts, S.-C. 125, 141

Batelaan, J.G. 81
Bertram, D. 219
Bhattacharjee, A.K. 271
Bliard, C. C1
Bollhagen, R. 113
Botter, H. 81

Castillón, S. C1
Ćetković, G. 261
Cherniak, R. 205
Clayden, N.J. 1
Cleophax, J. C5
Corrêa, J.B.C. 265
Csanádi, J. 251

Dazzo, F.B. 151
De Bruin, A.H. 195
Djurendić, E. 251
Dobson, C.M. 1
Dowd, M.K. 15
Dubreuil, D. C5

Efremenko, V.I. 185
El-Laghdach, A. C1

Farber, S.M. 185
French, A.D. 15

Gaši, K. 251
Gero, S.D. C5
Glaudemans, C.P.J. 101, 271
Gómez-Guillén, M. 255
Gómez-Sánchez, A. 255
Goux, W.J. 65
Gruppen, H. 45

Hasegawa, A. 87
Heyes, S.J. 1
Higuchi, R. 231

Hoffmann, R.A. 45
Hollingsworth, R.I. 151

Isobe, R. 231

Kamerling, J.P. 45, 81
Kiso, M. 87
Knirel, Yu.A. 185
Kochetkov, N.K. 185
Köll, P. 35
Komori, T. 231
Kopf, J. 35
Kormelink, F.J.M. 45
Kováč, P. 101
Kragten, E.A. 81
Kwon-Chung, K.J. 205, 271

Lajšić, S. 261
Legler, G. 113
Lehmann, J. 125, 141
Luger, P. 225
Lukacs, G. C1

Machado, A.S. C5
Martín-Zamora, M.-E. 255
Matheu, M.I. C1
Matoub, M. 273
Matta, K.L. 245
Mazid, M.A. 175
Messner, P. 175
Miljković, D. 251, 261
Mora, P. 273
Morf, M. 35
Mukasa, H. 237
Müller, R. 219
Murdoch, K.A. 161

Olesker, A. C1
Orgambide, G.G. 151

Paramonov, N.A. 185
Parolis, H. 195
Parolis, L.A.S. 195

Petek, F. 273
Petráková, E. 101
Petry, S. 125, 141
Pommerening, K. 219
Prabhanjan, H. 87

Rabius, E. 225
Reicher, F. 265
Reilly, P.J. 15
Rein, H. 219
Reiss, E. 205
Rouland, C. 273

Sarkar, A.K. 245
Shashkov, A.S. 185
Shimamura, A. 237
Sierakowski, M.-R. 265
Sleytr, U.B. 175

Thomas, N.F. C5
Tsumori, H. 237
Turner, S.H. 205

Uezono, Y. 237
Unger, F.M. 175

Vliegenthart, J.F.G. 45, 81
Voragen, A.G.J. 45
Vukojević, N. 251

Weber, D.S. 65

Yarullin, R.G. 185

Zawadzki-Baggio, S.F. 265
Zimmer, B. 35

SUBJECT INDEX

- (\pm)-6-Acetamido-1,2-anhydro-6-deoxy-*myo*-inositol, a tight-binding inhibitor and pseudo-substrate for *N*-acetyl- β -glucosaminidases, 113
- N*-Acetyl- β -glucosaminidases, studies of (\pm)-6-acetamido-1,2-anhydro-6-deoxy-*myo*-inositol as a tight-binding inhibitor and pseudosubstrate for, 113
- 4-(Adenin-9-yl)-1,6-anhydro-4-deoxy- β -D-glucopyranose dihydrate, structure of, 225
- Amicetose, an improved synthesis of D-, 261
- Amylose-iodine complex, studies of the nature of the, 161
- 1,6-Anhydro-4-deoxy- β -D-glucopyranose dihydrate, (4-adenin-9-yl)-, structure of, 225
- Anion-exchange chromatography with pulsed amperometric detection, composition analysis of sulfoethylcelluloses by high-pH, 81
- Antigens, artificial. Synthetic carbohydrate hap-
tens immobilized on crystalline bacterial surface layer glycoproteins, 175
- α -L-Arabinofuranan, linear (1 \rightarrow 5)-linked, from the seeds of guapuruvu (*Schizolobium parahybum*), 265
- Arabinoxylan, characterisation by ^1H NMR spectroscopy of enzymically derived oligosaccharides from alkali-extractable wheat-flour, 45
- Assignment of ^{13}C NMR signals for reduced nigerooligosaccharides prepared by partial acid hydrolysis of (1 \rightarrow 3)- α -D-glucan, 237
- Bead-cellulose, estimation of dialdehyde groups in 2,3-dialdehyde, 219
- Capsular polysaccharide of *Cryptococcus neoformans* serotype B (clarification), 271
- Capsular polysaccharide of *E. coli* serotype K47, structural elucidation, 195
- Carbohydrates, expedient syntheses of inososes by the Ferrier reaction of, C5
- Cellobiose, sophorose, and laminarabiose, conformational analysis of the anomeric forms of, using MM3, 15
- Cellulose, estimation of dialdehyde groups in 2,3-dialdehyde bead-, 219
- Chelating properties, a derivative of chitosan and 2,4-pentanedione with strong, 255
- Chitosan and 2,4-pentanedione, a derivative with strong chelating properties, 255
- Composition analysis of sulfoethylcelluloses by high-pH anion-exchange chromatography with pulsed amperometric detection, 81
- Conformational analysis of anomeric forms of sophorose, laminarabiose, and cellobiose using MM3, 15
- ^{13}C -CP/MAS NMR studies of the cyclomalto-oligosaccharide (cyclodextrin) hydrates, 1
- Cryptococcus neoformans* serotype A, structural variability in the glucuronoxylomannan of, determined by ^{13}C NMR spectroscopy, 205
- Cryptococcus neoformans* serotype B, major capsular polysaccharide of (clarification), 271
- Crystal and molecular structures of the pentitol penta-acetates, 35
- Cyclomalto-oligosaccharide (cyclodextrin) hydrates, ^{13}C -CP/MAS NMR studies, 1
- 2-Deoxy-2,2-difluoro-D-glucose (2-deoxy-2,2-difluoro-D-*arabino*-hexose), a new and extremely fast synthesis, C1
- 2-Deoxy-2-iodo-D-mannopyranose derivatives, a novel route to, 251
- Deoxygenated methyl α -isomaltotriosides, syntheses, 101
- Dialdehyde groups in 2,3-dialdehyde bead-cellulose, estimation of, 219
- Disialoganglioside GD1 α and its positional isomer, synthesis, 87
- E. coli* serotype K47, structural elucidation of the capsular polysaccharide of, 195

- Enzymically derived oligosaccharides from alkali-extractable wheat-flour arabinoxylan, characterisation by ^1H NMR spectroscopy of, 45
- Ferrier reaction, conformational and stereoelectronic aspects of the, C5
- β -D-Galactopyranosyl, 2,3,4,6-tetra-*O*-benzyl-, phenyl sulfoxide, use as a glycosylating donor of, 245
- α -D-Galactopyranosyl group, synthesis of oligosaccharides containing a, 245
- (1 \rightarrow 4)- β -D-Galactosyltransferase, a spacer-modified disaccharide as a photoaffinity reagent for the acceptor-binding area of bovine, 125
- (1 \rightarrow 4)- β -D-Galactosyltransferase, spacer-modified trisaccharide glycosides that mimic the biantennary Asn-linked oligosaccharide acceptor of, 141
- Ganglioside GD1 α and a positional isomer, synthesis, 87
- Gangliosides, negative-ion, fast-atom bombardment mass spectrometry of, 231
- β -D-Glucopyranose dihydrate, 4-(adenin-9-yl)-1,6-anhydro-4-deoxy-, structure of, 225
- Glucose, a new and extremely fast synthesis of 2-deoxy-2,2-difluoro-D-, C1
- β -Glucosidases purified from the termite *Macrotermes muelleri* and from its symbiotic fungus *Termitomyces* sp., properties, 273
- Glucuronoxylomannan of *Cryptococcus neoformans* serotype A, structural variability in the, determined by ^{13}C NMR spectroscopy, 205
- Glycolipid from *Rhizobium trifoli* ANU843, structural characterization of a novel diglycosyl diacylglyceride, 151
- Guapuruvu, linear (1 \rightarrow 5)-linked α -L-arabinofuranan from the seeds of, 265
- High-polar matrix system, negative-ion, fast-atom bombardment mass spectrometry of gangliosides using a, 231
- Inososes, expedient syntheses from carbohydrates, and conformational and stereoelectronic aspects of the Ferrier reaction, C5
- myo*-Inositol, (\pm)-6-acetamido-1,2-anhydro-6-deoxy-, a tight-binding inhibitor and pseudo-substrate for *N*-acetyl- β -glucosaminidases, 113
- Iodine, studies of the nature of the complex formed by amylose with, 161
- 2-Iodo-D-mannopyranose derivatives, a novel route to 2-deoxy-, 251
- α -Isomaltotriosides, methyl, syntheses of specifically deoxygenated, 101
- Laminarabiose, sophorose, and cellobiose, conformational analysis of the anomeric forms of, using MM3, 15
- Lipopolysaccharides, structure of the polysaccharide chains of *Pseudomonas pseudomallei*, 185
- D-Mannopyranose derivatives, a novel route to 2-deoxy-2-iodo-, 251
- Mass spectrometry, negative-ion, fast-atom bombardment, of gangliosides, 231
- MM3, conformational analysis of anomeric forms of sophorose, laminarabiose, and cellobiose using, 15
- Molecular and crystal structures of the pentitol penta-acetates, 35
- Nigeroooligosaccharides prepared by partial acid hydrolysis of (1 \rightarrow 3)- α -D-glucan, assignment of ^{13}C NMR signals for reduced, 237
- NMR pattern-recognition methods to the classification of peracetylated oligosaccharide residues, application of: effects of intraclass structure, 65
- NMR signals for reduced nigeroooligosaccharides prepared by partial acid hydrolysis of (1 \rightarrow 3)- α -D-glucan, assignment of ^{13}C , 237
- NMR studies of the cyclomalto-oligosaccharide (cyclodextrin) hydrates, ^{13}C -CP/MAS, 1
- Oligosaccharide acceptor of (1 \rightarrow 4)- β -D-galactosyltransferase, spacer-modified trisaccharide glycosides that mimic the biantennary Asn-linked, 141
- Oligosaccharides from alkali-extractable wheat-flour arabinoxylan, characterisation by ^1H NMR spectroscopy of enzymically derived, 45
- 2,4-Pentanedione and chitosan, a derivative with strong chelating properties, 255
- Pentitol penta-acetates, the crystal and molecular structure of the, 35
- Photoaffinity reagent for the acceptor-binding area of bovine (1 \rightarrow 4)- β -D-galactosyltransferase, a spacer-modified disaccharide as a, 125
- Polysaccharide chains of *Pseudomonas pseudomallei* lipopolysaccharides, structure of, 185
- Polysaccharide of *E. coli* serotype K47, structural elucidation of the capsular, 195
- Pseudomonas pseudomallei* lipopolysaccharides, structure of the polysaccharide chains of, 185

Rhizobium trifoli ANU843, structural characterization of a novel diglycosyl diacylglyceride glycolipid from, 151

Sophorose, laminarabiose, and cellobiose, conformational analysis of the anomeric forms of, using MM3, 15

Spacer-modified disaccharide as a photoaffinity reagent for the acceptor-binding area of bovine (1 → 4)- β -D-galactosyltransferase, 125

Spacer-modified trisaccharide glycosides that mimic the biantennary Asn-linked oligosaccharide acceptor of (1 → 4)- β -D-galactosyltransferase, 141

Structural characterization of a novel diglycosyl

diacylglyceride glycolipid from *Rhizobium trifoli* ANU843, 151

Sulfoethylcelluloses, composition analysis by high-pH anion-exchange chromatography with pulsed amperometric detection, 81

Sulfoxide, tetra-*O*-benzyl- β -D-galactopyranosyl phenyl, use as a glycosylating donor of, 245

Synthesis of D-amicetose, an improved, 261

Synthesis of 2-deoxy-2,2-difluoro-D-arabinohexose (2-deoxy-2,2-difluoro-D-glucose), a new and extremely fast, C1

Termite *Macrotermes muelleri* and its symbiotic fungus *Termitomyces* sp., properties of two β -glucosidases from, 273

